

REFERENCES:

1. Alcoa of Australia (2002). Indicators of *P. cinnamomi* used by Interpreters. Environment Research Bulletin.
2. Barrett, S., Shearer, B. L., Crane, C. E., & Cochrane, A. (2008). An extinction-risk assessment tool for flora threatened by *Phytophthora cinnamomi*. *Australian Journal of Botany*, 56; 477-486.
3. Bluett, V., Weste, G., & Cahill, D. (2003). Distribution of disease caused by *Phytophthora cinnamomi* in Wilsons Promontary National Park and potential for further impact. *Australasian Plant Pathology*, 32; 479-491.
4. Cahill, D. M., Rookes, J. E., Wilson, B. A., Gibson, L., & McDougall, K. L. (2008). Turner Review no. 17: *Phytophthora cinnamomi* and Australia's biodiversity: impacts, predictions and progress towards control. *Australian Journal of Botany*, 56; 279-310.
5. Cho, J. J. (1983). Variability in susceptibility of some *Banksia* Species to *Phytophthora cinnamomi* and their distribution in Australia. *Plant Disease*, 67; 869-871.
6. D'Souza, N. K., Colquhoun, I. J., Shearer, B. L., & Hardy, G. E. S. J. (2002). Assessing the potential for biological control of *Phytophthora cinnamomi* by fifteen native Western Australian jarrah-forest legume species. *Australasian Plant Pathology*, 34; 533-540.
7. Edmiston, R. J. (1989). Plants Resistant to Dieback. Department of Conservation and Land Management. 1-89.
8. Environment Australia. (2001). Threat Abatement Plan for Dieback caused by the root-rot fungus *Phytophthora cinnamomi*.
9. Gardner, J. H. & Rokich, P. A. (1987). *Phytophthora cinnamomi* in operational and rehabilitated bauxite mine area in South-western Australia. Alcoa of Australia Environment Research Bulletin. 13.
10. Hart, R. (1983). Report on Dieback due to *Phytophthora cinnamomi* in Two People's Bay Nature Reserve. Department of Fisheries and Wildlife.
11. Hill. (1990). Dieback Diseases and other *Phytophthora* spp. in the Northern Kwongan. In *Nature, Conservation, Landscape and Recreational Values of the Lesueur Area. A report to the Environmental Protection Authority from the Department of Conservation and Land Management*. eds. A. Burbidge, S. D. Hopper & S. van Leeuwen, EPA. WA.
12. Keighery, G. (1992). The impact of *Phytophthora* species on rare plants. In *Dieback-What is the Future?*, eds. M. J. Freeman, R. Hart & M. Ryall, Northern Sandplains Dieback Working Group. Perth, pp. 29-36.
13. Laidlaw, W. S. & Wilson, B. A. (2003). Floristic and structural characteristics of a coastal heathland exhibiting symptoms of *Phytophthora cinnamomi* infestation in the eastern Otway Ranges, Victoria. *Australian Journal of Botany*, 51; 283-293.
14. Lee, T. C. & Wicks, T. J. (1977). *Phytophthora cinnamomi* in native vegetation in south Australia. *Australasian Plant Pathology Newsletter*, 6; 22-23.
15. Marks, G. C. & Smith, I. W. (1991). *The cinnamon fungus in Victorian forests*. Victorian Government Publications, Melbourne
16. McCredie, T. A., Dixon, K. W. & Sivasithamparam, K. (1985). Variability in the resistance of *Banksia* L.f. species to *Phytophthora cinnamomi*. *Australian Journal of Botany*, 33; 629-637.

17. McDougall, K. L., Hardy, G. E. S. J. & Hobbs, R. J. (2001). Additions to the host range of *Phytophthora cinnamomi* in the jarrah (*Eucalyptus marginata*) forest of Western Australia. *Australian Journal of Botany*, 49; 193-198.
18. McDougall, K. L., Hobbs, R. J. & Hardy, G. E. S. J. (2002). Vegetation of *Phytophthora cinnamomi*-infested and adjoining uninfested sites in the northern jarrah (*Eucalyptus marginata*) forest of Western Australia. *Australian Journal of Botany*, 50; 277-288.
19. McDougall, K. L., Hobbs, R. J. & Hardy, G. E. S. J. (2005). Distribution of understorey species in forest affected by *Phytophthora cinnamomi* in south-western Western Australia. *Australian Journal of Botany*, 53; 813-819.
20. McDougall, K. L., Summerell, B. A., Coburn, D., & Newton, M. (2003). *Phytophthora cinnamomi* causing disease in subalpine vegetation in New South Wales. *Australasian Plant Pathology*, 32; 113-115.
21. Panjehkeh, N., Backhouse, D., & Taji, A. (2007). Flower colour is associated with susceptibility to disease in the legume *Swainsona formosa*. *Australasian Plant Pathology*, 36; 341-346.
22. Podger, F. D. (1972). *Phytophthora cinnamomi*, a cause of lethal disease in indigenous plant communities in Western Australia. *Phytopathology*, 62; 972-981.
23. Reiter, N., Weste, G., & Guest, D. (2004). The risk of extinction resulting from disease caused by *Phytophthora cinnamomi* to endangered, vulnerable or rare plant species endemic to the Grampians, western Victoria. *Australian Journal of Botany*, 52; 425-433.
24. Shearer, B.L., Crane, C. E., Barrett, S., & Cochrane, A. (2007). Assessment of threatened flora susceptibility to *Phytophthora cinnamomi* by analysis of disease progress curves in shadehouse and natural environments. *Australasian Plant Pathology*, 36; 609-620.
25. Shearer, B.L., Crane, C. E., & Cochrane, A. (2004). Quantification of the susceptibility of the native flora of the Southwest Botanical Province, Western Australia, to *Phytophthora cinnamomi*. *Australian Journal of Botany*, 52; 435-443.
26. Shearer, B.L., Crane, C. E., & Cochrane, A. (2004). Phosphite reduces disease extension of a *Phytophthora cinnamomi* front in *Banksia* woodland, even after fire. *Australasian Plant Pathology*, 33; 249-254.
27. Shearer, B. L. & Dillon, M. (1996). Susceptibility of plant species in *Banksia* woodlands on the Swan Coastal Plain, Western Australia, to infection by *Phytophthora cinnamomi*. *Australian Journal of Botany*, 44; 433-445.
28. Shearer, B. L. & Dillon, M. (1995). Susceptibility of plant species in *Eucalyptus marginata* forest to infection by *Phytophthora cinnamomi*. *Australian Journal of Botany*, 43; 113-134.
29. Shearer, B. L., & Fairman, R. G. (2007). Application of phosphite in a high-volume foliar spray delays and reduces the rate of mortality of four *Banksia* species infected with *Phytophthora cinnamomi*. *Australasian Plant Pathology*, 36; 358-368.
30. Suddaby, T., Alhussaen, K., Daniel, R., & Guest, D. (2008). Phosphonate alters the defense responses of *Lambertia* species challenged by *Phytophthora cinnamomi*. *Australian Journal of Botany*, 56; 550-556.
31. Titze, J. F. & Palzer, C. R. (1969). Host list of *Phytophthora cinnamomi* Rands with special reference to Western Australia. Forestry and Timber Bureau, Department of National Development. Technical Note No. 1.
32. Tynan, K. M., Scott, E. S. & Sedgley, M. (1998). Evaluation of *Banksia* species for response to *Phytophthora* infection. *Plant Pathology*, 47; 446-455.

33. Tynan, K. M., Wilkinson, C. J., Holmes, J. M., Dell, B., Colquhoun, I. J., McComb, J. A. & Hardy, G. E. St. J. (2001). The long-term ability of phosphite to control *Phytophthora cinnamomi* in two native plant communities of Western Australia. *Australian Journal of Botany*, 49; 761-770.
34. Weste, G. (2003). The dieback cycle in Victorian forests: a 30-year study of changes caused by *Phytophthora cinnamomi* in Victorian open forests, woodlands and heathlands. *Australasian Plant Pathology*, 32; 247-256.
35. Weste, G., Brown, K., Kennedy, J., & Walshe, T. (2002). *Phytophthora cinnamomi* infestation - a 24-year study of vegetation change in forests and woodlands of the Grampians, Western Victoria. *Australian Journal of Botany*, 50; 247-274.
36. Wills, R. T. (1993). Unpublished data.